

Spin Stiffness of Mesoscopic Quantum Antiferromagnets

Daniel Loss¹ and Dmitrii L. Maslov^{1,2*}

¹ *Department of Physics, Simon Fraser University, Burnaby B.C. V5A 1S6, Canada*

² *Department of Physics and Materials Research Laboratory, University of Illinois at Urbana-Champaign, IL 61801*

We study the spin stiffness of a one-dimensional quantum antiferromagnet in the whole range of system sizes L and temperatures T . We show that for integer and half-odd integer spin case the stiffness differs fundamentally in its L and T dependence, and that in the latter case the stiffness exhibits a striking dependence on the parity of the number of sites. Integer spin chains are treated in terms of the non-linear sigma model, while half-odd integer spin chains are discussed in a renormalization group approach leading to a Luttinger liquid with Aharonov-Bohm type boundary conditions.

PACS numbers: 75.10.Jm, 75.50.Ee, 73.20.Fz

Quantum one-dimensional antiferromagnets have been the subject of intensive studies since Haldane [1] conjectured that the spectrum of an integer spin S chain has a finite gap even in the absence of any anisotropy, while half-odd integer S chains are gapless. In both cases the Néel long-range order of the ground state is destroyed by quantum fluctuations. However, the “degree of destruction” is different: for integer S the correlation length is finite, which means that the elementary excitations have a gap, while for half-odd integer S the correlation length is infinite and excitations are gapless. By now, the presence of the Haldane gap for integer S chains is well understood theoretically [2] and has been confirmed in experimental [3] and numerical [4] studies. However, all these investigations were concentrated on the Haldane gap, i.e., on the energy spectrum itself. Thus a broader understanding of the Haldane conjecture is desirable, and the question naturally arises whether there are alternative manifestations of the fundamental difference between integer and half-odd integer spin chains. Indeed, it is the purpose of this work to provide an affirmative answer to this question and to discuss such a particular case in terms of the so-called spin stiffness.

Quite recently, there has been much interest in the spin stiffness (helicity modulus) ρ_s of *classical* Heisenberg ferromagnets [5–7]. ρ_s is defined as a change in the free energy F of the magnet when a twist is applied to the spins at the sample boundaries. When thermal fluctuations are taken into account, ρ_s is being renormalized with respect to its bare value and depends on the scale at which it is probed. Chakravarty [5] has recently shown that ρ_s exhibits features which are familiar from the behavior of the electrical conductance of a metal in the weak localization regime [8]. For instance, in 2d, the mean value of ρ_s depends logarithmically on the sample size L , while the rms value of its fluctuations is universal [9]. This similarity [10] makes the spin stiffness an equal member of the family of traditionally mesoscopic quantities such as a conductance or a persistent current. Also, the stiffness ρ_s serves as a useful (though not perfect) tool for characterizing magnetic long-range order; in particular, a vanishing value of ρ_s indicates absence of order [11].

Our goal here is to study ρ_s of a *quantum* one-dimensional antiferromagnet, where fluc-

tuations are i) quantum and ii) topologically distinct for integer and half-odd integer S . We shall see that the behavior of ρ_s is, indeed, quite different for integer and half-odd integer S : ρ_s is renormalized with L in the former case (as it is for a classical 2d ferromagnet), whereas it is L -independent (in leading order) in the latter. Moreover, for half-odd integer S , ρ_s is shown to exhibit a striking dependence on the parity of the total number N of spins. These results should be amenable to a direct check in numerical simulations (see e.g. Refs. [7,12]), and, in particular, could be tested experimentally by measuring the stiffness of quasi-one-dimensional antiferromagnets of finite size using similar materials as in Ref. [3].

We start with the Heisenberg Hamiltonian for a spin chain with nearest-neighbor interactions

$$H = J_{ex} \sum_{n=1}^N \mathbf{S}(n) \mathbf{S}(n+1), \quad (1)$$

where $J_{ex} > 0$, and we consider the integer S case first. In this case, the long-wavelength limit of the partition function Z becomes the $(1+1)$ d non-linear σ model (NL σ M) [13] with $Z = \int \mathcal{D}\mathbf{n} \delta(\mathbf{n}^2 - 1) \exp(-\mathcal{A})$, and the Euclidean action is given by

$$\mathcal{A} = \frac{1}{2g} \int_0^{L_T} \int_0^L dx_0 dx (\partial_\mu \mathbf{n})^2, \quad \mu = 0, 1, \quad (2)$$

where $g = 2/S$, $v_s = 2SJ_{ex}a_0$ is the spin wave velocity (we set $k_B = \hbar = 1$), a_0 is the lattice constant, $L = a_0N$, $L_T = \beta v_s$ is the wavelength of the thermal magnons, and $\mathbf{n}$ is the slow-varying component of the (staggered) magnetization satisfying the constraint $\mathbf{n}^2 = 1$. On the edges of the space-time domain $L \times L_T$ the boundary conditions are periodic in x_0 , i.e., $\mathbf{n}(0, x) = \mathbf{n}(L_T, x)$, and correspond to a fixed twist of the $\mathbf{n}$ -field applied in x -direction, i.e., $\mathbf{n}(x_0, 0) = (1, 0, 0)$ and $\mathbf{n}(x_0, L) = (\cos \theta, \sin \theta, 0)$, where θ is the twist angle. It is convenient to use the transformation [5,6,14] $\mathbf{n} = \mathcal{R}(\theta(x))\sigma$, where $\mathcal{R}$ is the rotation matrix about the z -axis by the angle $\theta(x) = \theta x/L$, and σ satisfies the boundary conditions $\sigma_1 = 1$, and $\sigma_{2,3} = 0$, at $x = 0, L$. The action then takes the form

$$\begin{aligned} \mathcal{A} = \frac{1}{2g} \int d\mathbf{x} [(\partial_\mu \sigma)^2 + \frac{\theta^2}{L^2} (1 - \sigma_3^2) \\ + 2 \frac{\theta}{L} (\sigma_1 \partial_x \sigma_2 - \sigma_2 \partial_x \sigma_1)]. \end{aligned} \quad (3)$$

We define the spin stiffness ρ_s in units of the velocity

$$\rho_s = \frac{1}{2}L \frac{\partial^2 F}{\partial \theta^2} \Big|_{\theta=0}, \quad (4)$$

where $F = -T \ln Z$ [15]. With this definition, the bare value of ρ_s in the tree (classical) approximation is $\rho_s^0 = v_s/2g$. Corrections due to quantum and thermal fluctuations can be found in a loop expansion in g . In one-loop order, only quadratic terms in the action (3) are to be retained, and the third term in Eq. (3) reduces to a total derivative $\partial_x \sigma_2$, and, thus, vanishes as we shall restrict our consideration to the topological sector with zero winding number (Pontryagin index). Performing the functional integration, we get

$$\rho_s = \rho_s^0 \left(1 - \frac{g}{LL_T} \sum_{\mathbf{q}} \frac{1}{\mathbf{q}^2} \right), \quad (5)$$

where $\mathbf{q} = (2\pi n/L_T, \pi m/L)$, $n = 0, \pm 1, \dots$, $m = 1, 2, \dots$. The sum in Eq. (5) can be evaluated in three limiting cases: a) $L \ll L_T$ (quantum region); b) $a_0 \ll L_T \ll L$ (classical renormalized region); and c) $L_T \ll a_0$ (classical region). We have

$$\frac{\rho_s}{v_s} = \begin{cases} \ln(\xi_{qm}/L)/4\pi, & \text{for a)} \\ (\xi(T) - L)/12L_T, & \text{for b)} \\ (\xi_{cl} - L)/12L_T, & \text{for c)} \end{cases} \quad (6)$$

Here, $\xi_{qm} = \alpha a_0 \exp(\pi S)$ is the correlation length in the quantum region. $\xi(T) = 3L_T \ln(\gamma \xi_{qm}/L_T)/\pi$ is the classical correlation length $\xi_{cl} = 6L_T/g$ renormalized by quantum fluctuations. α and γ are (cut-off dependent) non-universal constants of order one. We note that case c) agrees with the 1d classical NL σ M [5].

In all the regions, ρ_s goes to zero as the system size $L < \xi$ approaches the correlation length of the corresponding region [16]. This zero value of ρ_s is what a macroscopic system is expected to have in the absence of spontaneously broken symmetry [11] (the point $\rho_s = 0$ signals the breakdown of the one-loop order expansion).

The rms fluctuation $\delta \rho_s^2 = (TL \partial/\partial \theta^2)^2 \ln Z|_{\theta=0}$ is given by

$$\frac{\delta \rho_s^2}{v_s^2} = \frac{1}{2L^2 L_T^2} \sum_{\mathbf{q}} \frac{1}{\mathbf{q}^4} = \begin{cases} \frac{\zeta(3)}{8\pi^3} \frac{L}{L_T}, & \text{for a)} \\ \frac{1}{180} \left(\frac{L}{L_T} \right)^2, & \text{for b) and c),} \end{cases} \quad (7)$$

where $\zeta(x)$ is the Riemann ζ -function. Contrary to the case of a classical ferromagnet [5], the fluctuations are *non-universal*: they depend both on L and T . Moreover, in the classical and classical renormalized regions, the fluctuations are abnormally large ($\delta\rho_s > \rho_s$), and, thus, the spin stiffness is not a self-averaging quantity. Finally, we note that the analogues of the regions we consider in the quantum NL σ M can also be obtained in the classical 2d model, if one considers a rectangular instead of a square system.

We now turn to the half-odd integer spin case. The effective field-theoretical description of the long-wavelength excitations is not of much use here, since the partition function contains a Θ -term and contributions from all topological sectors with different winding numbers [1,2,13], which makes the model hardly tractable. It is believed that the exactly solvable case of the spin 1/2 chain reproduces the generic features of all half-odd integer S chains [1,2,13], and we shall consider this case only. By using the Jordan–Wigner transformation [17]

$$\psi(n) = (-1)^n \exp \left(i\pi \sum_{j=1}^{n-1} (S_z(j) + \frac{1}{2}) \right) S_-(n), \quad (8)$$

where $S_{\pm} = S_x \pm iS_y$, the Hamiltonian (1) is mapped on to a system of spinless fermions on the lattice,

$$H = J_{ex} \sum_{n=1}^N \left(-\frac{1}{2} \left\{ \psi^\dagger(n) \psi(n+1) + \text{H. c.} \right\} + : \rho(n) :: \rho(n+1) : \right), \quad (9)$$

where $: \rho(n) := \psi^\dagger(n) \psi(n) - 1/2$ is the (Fermi-ordered) density operator. We now have to specify the boundary condition for the fermionic operators ψ . The quantum generalization of the classical boundary conditions for the spin field $\mathbf{n}$, used in the NL σ M treatment of the integer S case, is $S_{\pm}(N+1) = e^{\pm i\theta} S_{\pm}(1)$, and $S_z(N+1) = S_z(1)$. The boundary condition for ψ then follows from Eq. (8),

$$\psi(N+1) = e^{i[\pi(N_F+N)-\theta]} \psi(1), \quad (10)$$

where the number of fermions is $N_F = N/2$, if N is even, and $N_F = (N+1)/2$, if N is odd. We have also used the fact that $\sum S_z(n) = 0(1/2)$ for even (odd) N . The problem

defined by Eqs. (9) and (10) is similar to that of spinless electrons on a ring threaded by an Aharonov-Bohm flux θ , with the difference that here the boundary conditions depend on the parity of N . This parity dependence will result in a striking difference in the behavior of ρ_s for even and odd N .

Finite-size systems of interacting fermions with twisted boundary conditions have recently been studied in the framework of the Luttinger liquid approach [18]. The parity dependence of the boundary conditions, however, requires a re-examination of the bosonization scheme, which we now address. The left- and right-movers are introduced by $\psi(n) = e^{ink_F}\psi_+(n) + e^{-ink_F}\psi_-(n)$, where we choose $k_F = \pi/2$ for N even and odd. The boundary conditions for ψ_α , where $\alpha = \pm$, take the form

$$\psi_\alpha(N+1) = \begin{cases} e^{-i\theta}\psi_\alpha(1), & \text{for } N \text{ even} \\ -\alpha ie^{-i\theta}\psi_\alpha(1), & \text{for } N \text{ odd.} \end{cases} \quad (11)$$

Bosonic fields are introduced by $\psi_\alpha = (2\pi a_0)^{-1/2} e^{i\sqrt{\pi}\phi_\alpha}$, where $\phi_\alpha = \alpha\phi - \vartheta$, and $\partial_x\vartheta$ is the conjugate momentum of ϕ . The zero modes of ϕ and ϑ can be chosen in the form [18]

$$\begin{aligned} \phi_0 &= \phi_J/\sqrt{\pi} + \mathbf{M}\sqrt{\pi}x/L, \\ \vartheta_0 &= \vartheta_M/\sqrt{\pi} + (\mathbf{J} - \theta/\pi)\sqrt{\pi}(x + \tfrac{1}{2}L)/L, \end{aligned} \quad (12)$$

where $\mathbf{J}$ and $\mathbf{M}$ are the operators of the topological current and the number of particles above the ground state [19], respectively, which satisfy $[\phi_J, \mathbf{J}] = [\vartheta_M, \mathbf{M}] = i$. Next, using the Baker-Hausdorff formula, we write

$$e^{i\sqrt{\pi}\phi_\alpha} = \bar{\psi} e^{i\frac{\pi}{L}[(\alpha x(M-1) - J(x + \frac{L}{2}))]} \quad (13)$$

where $\bar{\psi}$ contains contributions from the nonzero modes and from ϕ_J and ϑ_M , and is not parity dependent; $J(M)$ stands for the eigenvalue of $\mathbf{J}(\mathbf{M})$. It is convenient to introduce the topological indices κ_J and κ_M , such that $\kappa_J = 0(1)$, if J is even (odd); for even N , $\kappa_M = 0(1)$, if M is even (odd), and, for odd N , $\kappa_M = 0(1)$, if $M + 1/2$ is even (odd). By using Eqs. (11) and (13), we see that $\kappa_{J,M}$ must satisfy the following constraints: $\kappa_J = 1$,

$\kappa_M = 0$ (and vice versa), if N is even, and $\kappa_J = \kappa_M$, if N is odd. By comparing these constraints with the analogous constraints of the fermionic problem [18], we can now say that the ground state of our spin system is *paramagnetic* for N even and *diamagnetic* for N odd.

The rest of the bosonization procedure is identical to that of Ref. [18], and the bosonized Euclidean action takes the sine-Gordon form

$$\begin{aligned} \mathcal{A}_b = \int_0^{\beta v_0} \int_0^L dx \{ & K_0 (\partial_\mu \phi)^2 + (i/L) \sqrt{\pi} \theta_0 \partial_0 \phi \\ & - (g_0/a_0^2) \cos(4\sqrt{\pi}\phi) \}, \end{aligned} \quad (14)$$

where $v_0 = v_s(1+4/\pi)^{1/2}$, $K_0 = v_0/2v_s$, $g_0 = 1/8\pi^2 K_0$, and $\theta_0 = \kappa_J - \theta/\pi$. The bosonic fields have been decompactified in the course of the functional integral derivation, and ϕ obeys now the boundary conditions $\phi(x_0 + k_0\beta v_0, x + k_1L) = \phi(x_0, x) + k_0\sqrt{\pi}n + k_1\sqrt{\pi}(2m + \kappa_M)$, where n and m are the winding numbers in x_0 - and x -directions, respectively. The measure $\mathcal{D}\phi$ of the functional integral $Z = \int \mathcal{D}\phi \exp(-\mathcal{A}_b)$ includes the sums over the winding numbers n, m and over the topological indices $\kappa_{J,M}$. The last term in Eq. (14) corresponds to Umklapp scattering processes between fermions. Since g_0/K_0 is small (≈ 0.02), this Umklapp term can be treated perturbatively in a standard renormalization group (RG) approach leading to the following flow equations [20]:

$$\begin{aligned} \frac{dg}{dl} &= 2(K-1)g, \quad g(0) = g_0 \\ \frac{dK}{dl} &= 2\pi^2 g^2, \quad K(0) = K_0, \end{aligned} \quad (15)$$

where $l = \ln(\mathcal{L}/a_0)$ with $\mathcal{L} = \min\{L, L_T\}$. Since we started with the isotropic Heisenberg model (1), the scaling dimension of the Umklapp term is equal to the critical dimension of the model ($= 2$), i.e., this term is marginally relevant. In this case, the flow proceeds along the separatrix between massive and massless phase. On this line [21], the solutions to Eqs. (15) are with $\mathcal{L} \gg L_0$

$$K = K^* - \frac{1}{2 \ln \frac{\mathcal{L}}{L_0}}, \quad g = \frac{K^* - K}{\pi}, \quad (16)$$

where $K^* = K(\infty)$ is the fixed point value, and the (non-universal) cut-off L_0 depends on (K_0, g_0) and is of the order a_0 . At the fixed point ($\mathcal{L} \rightarrow \infty$), $g = 0$ and the action (14) renormalizes to that of a Luttinger liquid with a topological term ($\propto \partial_0 \phi$), and with parameters renormalized through interactions: $(K_0, v_0) \rightarrow (K^*, v^*)$. By comparing with the exact Bethe-ansatz solution [24], one gets $K^* = 1$ and $v^* = \pi v_s/2$.

We can now calculate the fixed-point value of the spin stiffness, ρ_s^* , and its finite size and finite temperature corrections. Upon integrating out the zero modes, the twist-dependent part of Z becomes

$$Z_\theta = \sum_{\kappa_J, \kappa_M} e^{-\kappa_M b} \theta_3(z_J, e^{-a}) \theta_3(z_M, e^{-4b}), \quad (17)$$

where $\theta_3(z, q) = \sum q^{n^2} e^{2inz}$ is the Jacobi θ function, $z_J = \pi\theta_0/2$, $a = \pi K^* L/L_T^*$, $b = \pi K^* L_T^*/L$, $z_M = 2i\kappa_M b$, and $L_T^* = \beta v^*$. The results for the spin stiffness take simple forms in the limiting cases of low and high temperatures. For $L \ll L_T^*$, we obtain from (4) and (17)

$$\rho_s^* = \begin{cases} -(v^*/8K^{*2})(L_T^*/L), & \text{for } N \text{ even} \\ v^*/4\pi K^*, & \text{for } N \text{ odd,} \end{cases} \quad (18)$$

while for $L \gg L_T^*$ we get

$$\rho_s^* = (-1)^{N+1} 2(L/L_T^*) e^{-\pi\chi L/L_T^*}, \quad (19)$$

where $\chi = K^* + 1/4K^*$. To obtain the value of ρ_s away from the fixed point, we go back to the full action (14) and replace (K_0, g_0) by (K, g) from Eq. (16), treating the deviations from the fixed-point values as perturbations [21–23]. In first order, the Umklapp term gives no contribution, while the perturbation in $K - K^*$ leads to:

$$\rho_s = \rho_s^* \times \begin{cases} 1 + 1/[\alpha_N \ln(L/L_0)], & \text{for } L \ll L_T^* \\ \exp(3L\pi/[8L_T^* \ln(T_0/T)]), & \text{for } L \gg L_T^*, \end{cases} \quad (20)$$

where $\alpha_N = 1$ for even N , and $\alpha_N = 2$ for odd N ; the cut-off T_0 is of the order of v^*/L_0 . The last equation is valid for $T \leq T^* < T_0$, where $L_T(T^*) = a_0$. The L - and

T -dependent corrections to ρ_s^* resulting from the perturbation of the fixed-point action by the marginally irrelevant operator are larger than those coming from the expansion of Eq. (17) in $(L/L_T^*)^{\pm 1}$ at the fixed point. In particular, the exponential dependence of ρ_s^* on K^* in the high-temperature regime results in a significant T -dependent renormalization. This renormalization may be conjectured to remain significant in the intermediate regime $L \simeq L_T^*$ as well. We also note that the Umklapp processes lead to the breakdown of the single-parameter scaling of ρ_s : the latter scales with L/L_T^* at the fixed-point but acquires additional L/L_0 and T/T_0 scalings away from the fixed point (for a rough estimate, $L_0 \simeq a_0$, and $T_0 \simeq J_{ex}$). This breakdown can be detected in numerical and real experiments. The fluctuations in ρ_s can be calculated along the same lines as for ρ_s itself, and, in marked contrast to the integer S case, turn out to be exponentially small for all L/L_T .

The negative value of ρ_s for even N simply reflects the fact that in this case the free energy has a maximum at $\theta = 0$, and an arbitrarily small twist drives the system out of this state. Analysing ρ_s at finite θ (and low temperatures) we can see that ρ_s vanishes at some $\theta^* \simeq L/L_T^* \ll 1$ and then remains positive for all θ , thus exhibiting a crossover from the paramagnetic to the diamagnetic regime. The parity effects in the spin stiffness are quite similar to that in persistent currents of electronic systems [25,26,18]; in particular, the result obtained above can be checked without approximations for the special case of the XY -model by mapping it on to the exactly solvable problem of free fermions [27].

This work was supported by the NSERC of Canada (D.L. and D.L.M.) and in part by the US NSF through Grant No. DMR-89-20538 (D.L.M.).

REFERENCES

- * Present address. On leave from: Institute for Microelectronics Technology, Chernogolovka, Moscow District, 142432, Russia.
- [1] F. D. M. Haldane, Phys. Lett. **93A**, 464 (1983); Phys. Rev. Lett. **50**, 1153 (1983).
- [2] I. Affleck, Nucl. Phys. **B265**, 409 (1986); J. Phys. Cond. Mat. **1**, 3047 (1989).
- [3] W. J. L. Buyers et al., Phys. Rev. Lett. **56**, 371 (1986); J. P. Renard et al., Europhys. Lett. **3**, 945 (1987).
- [4] R. Botet and R. Julien, Phys. Rev. **B27**, 613 (1983); M. P. Nightingale and H. W. J. Blöte, Phys. Rev. **B33**, 659 (1986); H. J. Schulz and T. A. L. Ziman, Phys. Rev. **B33**, 6545 (1986); M. Takahashi, Phys. Rev. Lett. **62**, 2313 (1989); S. R. White, Phys. Rev. Lett. **69**, 2863 (1992).
- [5] S. Chakravarty, Phys. Rev. Lett. **66**, 318 (1991).
- [6] E. Brézin et al., J. Stat. Phys. **70**, 583 (1993).
- [7] M. Caffarel, P. Azaria, B. Delamotte, and D. Mouhanna, Europhys. Lett. **26**, 493 (1994).
- [8] B. L. Al'tshuler, Pis'ma Zh. Eksp. Teor. Fiz. **41** 530 (1985) [Sov. Phys. JETP Lett. **41**, 648 (1985)]; P. A. Lee and A. D. Stone, Phys. Rev. Lett. **55**, 1622 (1985).
- [9] These results have been verified up to two-loop order [6] and in Monte-Carlo simulations [7].
- [10] This similarity comes from the close relation between spin and charge stiffness by virtue of the Jordan–Wigner transformation [17] and from the connection between conductivity and charge stiffness by virtue of Kohn's theorem (see W. Kohn, Phys. Rev. **133**, A171 (1964), and B.S. Shastry and B. Sutherland, Phys. Rev. Lett. **65**, 243 (1990)).
- [11] N. Goldenfeld, *Lectures on Phase Transitions and The Renormalization Group* (Addison-Wesley, Reading, MA, 1992).

- [12] J. Bonca, J.P. Rodriguez, J. Ferrer, and K.S. Bedell, Phys. Rev. B **50**, 3415 (1994).
- [13] E. Fradkin, *Field Theories of Condensed Matter Systems* (Addison-Wesley, Reading, MA, 1991).
- [14] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* 2nd ed. (Oxford Univ. Press, New York, 1993).
- [15] Although F is an even function in θ , it is not necessarily a function of θ^2 (as we will see in the half-odd integer spin case, F can be a function of $|\theta|$). Thus, definition (4) is more general than that used in previous works.
- [16] We note that a similar calculation in the $(2+1)$ d NL σ M gives exactly the same correlation lengths as found by S. Chakravarty, B. I. Halperin, and D. R. Nelson, Phys. Rev. B **39**, 2344 (1989), by solving the RG equations.
- [17] P. Jordan and E. Wigner, Z. Phys. **47**, 631 (1928).
- [18] D. Loss, Phys. Rev. Lett. **69**, 343 (1992).
- [19] F. D. M. Haldane, J. Phys. C **14**, 2585 (1981); Phys. Rev. Lett. **47**, 1840 (1981).
- [20] J. M. Kosterlitz and D. Thouless, J. Phys. C **6**, 1181 (1973); T. Ohta and D. Jasnow, Phys. Rev. B **20**, 139 (1979).
- [21] The phase below the separatrix corresponds to the anisotropic Heisenberg model with powerlaw behavior: $K_A^* - K = (\pi g)^2 / (2 - 2K_A^*) = (1/2)(L_0/\mathcal{L})^{4-4K_A^*}$, with $1/2 \leq K \leq K_A^* < 1$, and $K = 1/2$ corresponding to the XY limit. Thus in this case, we find $(\mathcal{L}/L_0)^{2-2K_A^*}$ instead of $\ln(\mathcal{L}/L_0)$ below in Eq. (20) in *leading* power.
- [22] In leading order, the flow of v is slower than that of K and g , and we still can use the fixed point value v^* .
- [23] For a similar treatment of the spin susceptibility in the framework of conformal field theory, see S. Eggert, I. Affleck, and M. Takahashi, Phys. Rev. Lett. **73**, 332 (1994).

- [24] R. Shankar, Int. J. Mod. Phys. B**4**, 2371 (1990).
- [25] N. Byers and C. N. Yang, Phys. Rev. Lett.**7**, 46 (1961).
- [26] A. J. Leggett, in *Granular Nanoelectronics*, edited by D. K. Ferry, J. R. Barker, and C. Jacoboni, NATO ASI Ser. B, Vol. 251 (Plenum, NY, 1991), p. 297.
- [27] Since the relation between the parity of N and the dia- or paramagnetic response might appear counterintuitive, we would like to recall that our boundary conditions for $\mathbf{S}$ stand for a *closed* ring with N sites and have the physical meaning of inverting (at $\theta = \pi$) the sign of the interaction between $\mathbf{S}(N)$ and $\mathbf{S}(1)$. As one can show, *open* chains where only the last spin is twisted behave according to one's expectation, i.e., even (odd) chains exhibit dia(para)-magnetic response.